

amer: Using **lme4** to fit Generalized Additive Mixed Models

Fabian Scheipl

August 31, 2009

Abstract

The **lme4** package uses sparse matrix technology and clever likelihood-decompositions to fit linear, generalized, and nonlinear mixed-effects models. The **amer** package extends **lme4**'s scope to include generalized *additive* mixed models (GAMM). This vignette summarizes the main ideas behind additive models and their representation in the form of a mixed model, describes the modifications to **lmer** necessary for fitting GAMMs and presents some examples with real data.

Contents

1	Additive Models	3
2	Mixed model representation of an additive model	5
2.1	Variability Estimation	7
3	Examples	8
3.1	Generalized Additive Model	8
3.2	Separate smooths for levels of a factor: Using the <code>by</code> -option . .	11
3.3	Subject- or cluster-specific smooths: Using the <code>allPen</code> -option	15
3.4	Varying coefficient models: Using the <code>varying</code> -option	18
3.5	Variability Bands: Using the <code>RW</code> - and <code>MCMC</code> -options	21
4	Implementation	24
4.1	What's the point?	24
4.2	Making <code>lmer</code> fit GAMMs	24
4.3	Justification for using the TP-Basis	26
4.4	Writing your own basis-generating function	26
5	Open Issues	30

1 Additive Models

In many applications, the assumption of a linear dependence of the response on predictor variables is inappropriate. Modelling smooth functions of an unknown shape, that is, models of the form

$$\mathbf{y} = \sum_{s=1}^S f_s(\mathbf{x}_s) + \boldsymbol{\varepsilon}; \quad \boldsymbol{\varepsilon} \sim N_n(0, \sigma_\varepsilon^2 \mathbf{I}_n)$$

where $f_s(\cdot)$ is some smooth function of a covariate \mathbf{x}_i , which can also be multidimensional (e.g. surface estimation), requires solving 3 problems not encountered in linear modelling:

1. the smooth function has to be represented somehow
2. the degree of smoothness of the function must be controllable
3. the amount of smoothness most appropriate should be selected in a data-driven way

Spline smoothing addresses the first issue by assuming that $f_s(\mathbf{x}_s)$ can be approximated by a linear combination of d_s basis functions $B_j(\mathbf{x}_s)$, $j = 1, \dots, d_s$:

$$f_s(\mathbf{x}_s) \approx \mathbf{B}_s \boldsymbol{\delta}_s; \quad \mathbf{B}_s = \begin{bmatrix} B_1(x_{s1}) & \dots & B_{d_s}(x_{s1}) \\ \vdots & & \vdots \\ B_1(x_{sn}) & \dots & B_{d_s}(x_{sn}) \end{bmatrix}$$

This obviously leads back to a linear modelling context. For ease of notation we set $S = 1$ and drop the subscript s in the following.

The second issue, controlling the roughness or “wiggleness” of the estimated function, is a variant of the bias-variance tradeoff problem: using too few basis functions may not allow the fitted curve to accurately represent the shape of the function, leading to biased estimation, while using too many will result in an overly close interpolation of the measured data points — the estimated curve represents random noise along with the underlying structure. Penalized spline smoothing (Eilers and Marx, 1996) addresses this problem

by choosing a sufficient number of knots (e.g. 10-40) to ensure the necessary flexibility of the fit and by introducing an additional penalty term, a function of the spline coefficients $\boldsymbol{\delta}$, that quantifies the roughness of the estimated function. For a broad class of spline bases, the resulting criterion is a penalized least squares criterion,

$$\min_{\boldsymbol{\delta}} \left(\|\mathbf{y} - \mathbf{B}\boldsymbol{\delta}\|^2 + \frac{1}{\lambda} \boldsymbol{\delta}' \mathbf{K} \boldsymbol{\delta} \right), \quad (1)$$

where \mathbf{K} is a penalty matrix and λ is the smoothing parameter controlling the amount of penalization, i.e. the tradeoff between fidelity to the data and complexity of the fit. The elements in \mathbf{K} are determined by the spline basis that is used to generate \mathbf{B} and the roughness penalty desired by the analyst (usually penalizing (local) deviations of the fitted function from a constant, linear, or a quadratic polynomial).

Example: TP-Basis

A simple example of basis functions is the truncated powers (TP) basis. A TP-Basis of degree p , with d basis functions for a covariate \mathbf{x} and fixed knots $\kappa_1, \dots, \kappa_{d-p}$ consists of a constant term, p global polynomial terms $\mathbf{x}^1, \dots, \mathbf{x}^p$ and $p-d$ truncated polynomials $(\mathbf{x} - \kappa_i)_+^p$, $i = 1, \dots, d-p$, where $(y)_+ = y I(y > 0)$:

$$\mathbf{B} = \begin{bmatrix} \mathbf{x}^0 & \mathbf{x}^1 & \dots & \mathbf{x}^p & (\mathbf{x} - \kappa_1)_+^p & \dots & (\mathbf{x} - \kappa_{d-p})_+^p \end{bmatrix}$$

The penalty for the TP-Basis penalizes deviations of the fitted function from a p -degree polynomial:

$$\mathbf{K} = \text{diag}(\mathbf{0}_{p+1}, \mathbf{1}_{d-p}).$$

The penalty term $\boldsymbol{\delta}' \mathbf{K} \boldsymbol{\delta}$ is simply the sum of squares of the $p-d$ coefficients for the truncated polynomials.

2 Mixed model representation of an additive model

The third issue, selecting the amount of smoothness most appropriate in a data-driven way, then reduces to estimation of the smoothing parameter λ , which controls the smoothness of the estimated function. The penalized least squares problem is reformulated as a mixed model in which the smoothing parameter becomes a variance component. This is achieved by a decomposition of the spline coefficients into an unpenalized part and a penalized part:

$$\boldsymbol{\delta} = \boldsymbol{U}\boldsymbol{\beta} + \boldsymbol{P}\boldsymbol{b}$$

where \boldsymbol{U} , $d \times p$, is a basis of the p -dimensional nullspace of the penalization matrix \boldsymbol{K} and \boldsymbol{U} and \boldsymbol{P} have the following properties (Kneib, 2006, ch. 5.1):

1. The composed matrix $[\boldsymbol{U}\boldsymbol{P}]$ has full rank to make the transformation above a one-to-one transformation. This also implies that both \boldsymbol{U} and \boldsymbol{P} have full column rank.
2. \boldsymbol{U} and \boldsymbol{P} are orthogonal, i. e. $\boldsymbol{U}\boldsymbol{P}' = \mathbf{0}$
3. $\boldsymbol{U}'\boldsymbol{K}\boldsymbol{U} = \mathbf{0}$, so that $\boldsymbol{\beta}$ is unpenalized by \boldsymbol{K}
4. $\boldsymbol{P}'\boldsymbol{K}\boldsymbol{P} = \boldsymbol{I}$, so that the penalty for \boldsymbol{b} reduces to $\|\boldsymbol{b}\|^2$

The decomposition is not unique. It can always be based on the spectral decomposition of \boldsymbol{K} . With

$$\boldsymbol{K} = [\boldsymbol{\Lambda}_+ \boldsymbol{\Lambda}_0]' \begin{bmatrix} \boldsymbol{\Gamma}_+ & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} [\boldsymbol{\Lambda}_+ \boldsymbol{\Lambda}_0],$$

where $\boldsymbol{\Lambda}_+$ is the matrix of the eigenvectors associated with the positive eigenvalues $\text{diag}(\boldsymbol{\Gamma}_+)$, and $\boldsymbol{\Lambda}_0$ are the eigenvectors associated with the zero eigenvalues, the decomposition is

$$\begin{aligned} \boldsymbol{U} &= \boldsymbol{\Lambda}_0, \\ \boldsymbol{P} &= \boldsymbol{L}(\boldsymbol{L}'\boldsymbol{L})^{-1} \end{aligned}$$

with $\boldsymbol{L} = \boldsymbol{\Lambda}_+ \boldsymbol{\Gamma}_+^{1/2}$.

Using

$$\begin{aligned}
\mathbf{B}\boldsymbol{\delta} &= \mathbf{B}(\mathbf{U}\boldsymbol{\beta} + \mathbf{P}\mathbf{b}) = \mathbf{X}_u\boldsymbol{\beta} + \mathbf{Z}_p\mathbf{b} \\
\text{and } \boldsymbol{\delta}'\mathbf{K}\boldsymbol{\delta} &= (\mathbf{U}\boldsymbol{\beta} + \mathbf{P}\mathbf{b})'\mathbf{K}(\mathbf{U}\boldsymbol{\beta} + \mathbf{P}\mathbf{b}) \\
&= \mathbf{b}'\mathbf{P}'\mathbf{K}\mathbf{P}\mathbf{b} + \boldsymbol{\beta}'\mathbf{U}'\mathbf{K}\mathbf{U}\boldsymbol{\beta} = \mathbf{b}'\mathbf{b},
\end{aligned} \tag{2}$$

the penalized least squares criterion (1) can be rewritten as

$$\begin{aligned}
\min_{\boldsymbol{\delta}} \left(\|\mathbf{y} - \mathbf{B}\boldsymbol{\delta}\|^2 + \frac{1}{\lambda} \boldsymbol{\delta}'\mathbf{K}\boldsymbol{\delta} \right) &= \\
\min_{\boldsymbol{\beta}, \mathbf{b}} \left(\|\mathbf{y} - \mathbf{X}_u\boldsymbol{\beta} + \mathbf{Z}_p\mathbf{b}\|^2 + \frac{1}{\lambda} \|\mathbf{b}\|^2 \right). &
\end{aligned} \tag{3}$$

Example: TP-Basis

Since the penalty matrix $\mathbf{K} = \text{diag}(\mathbf{0}_{p+1}, \mathbf{1}_{d-p})$ of the TP-Basis has such an easy structure, the decomposition into the unpenalized part and the penalized part of the spline coefficients $\boldsymbol{\delta}$ is trivial: \mathbf{X}_u contains the first $p+1$ columns of \mathbf{B} with the constant term and the polynomials, while \mathbf{Z}_p contains the $d-p$ columns with the truncated polynomial terms:

$$\begin{aligned}
\mathbf{B} &= [\mathbf{X}_u \mathbf{Z}_p] \\
\mathbf{X}_u &= \begin{bmatrix} \mathbf{x}^0 & \mathbf{x}^1 & \dots & \mathbf{x}^p \end{bmatrix} \\
\mathbf{Z}_p &= \begin{bmatrix} (\mathbf{x} - \kappa_1)_+^p & \dots & (\mathbf{x} - \kappa_{d-p})_+^p \end{bmatrix}
\end{aligned}$$

For given λ , minimizing (3) over $(\boldsymbol{\beta}', \mathbf{b}')'$ is equivalent to BLUP-estimation (Ruppert et al., 2003, ch. 4.5.3) in a linear mixed model with

$$\mathbf{y} = \mathbf{X}_u\boldsymbol{\beta} + \mathbf{Z}_p\mathbf{b} + \boldsymbol{\varepsilon}; \quad \boldsymbol{\varepsilon} \sim N_n(0, \sigma_\varepsilon^2 \mathbf{I}_n); \quad \mathbf{b} \sim N_{d-p}(0, \sigma_\varepsilon^2 \lambda \mathbf{I}_{d-p}).$$

To see this, multiply the expression in (3) with $\frac{1}{2\sigma_\varepsilon^2}$: For given λ and σ_ε^2 , it is equivalent to maximizing

$$L(\boldsymbol{\beta}, \mathbf{b} | \lambda, \sigma_\varepsilon^2) \propto \exp \left(\frac{\|\mathbf{y} - \mathbf{X}_u\boldsymbol{\beta} - \mathbf{Z}_p\mathbf{b}\|^2 + \frac{1}{\lambda} \|\mathbf{b}\|^2}{-2\sigma_\varepsilon^2} \right),$$

which is the likelihood of the model above. In effect, the reformulation as a mixed model above allows for the estimation of smoothing parameters with

ML- or REML-methodology. All this applies as well for the generalized case with non-gaussian responses.

This model formulation can be extended to include multiple smooth terms, other random effects and a linear predictor in the classical sense of linear regression: Just concatenate the unpenalized parts of the smooth terms to the design matrix of the fixed effects and the penalized parts of the smooth effects to the design matrix of the random effects.

For a mixed additive model

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \sum_{l=1}^L \mathbf{Z}_l \mathbf{b}_l + \sum_{s=1}^S f(\mathbf{x}_s) + \varepsilon; \quad \begin{bmatrix} \mathbf{b} \\ \boldsymbol{\varepsilon} \end{bmatrix} \sim N_{\sum q_l + n} \left(\mathbf{0}, \sigma_\varepsilon^2 \begin{bmatrix} \boldsymbol{\Omega}_b^{-1} & 0 \\ 0 & \mathbf{I}_n \end{bmatrix} \right)$$

with fixed effects design \mathbf{X} , L random effects designs \mathbf{Z}_l each with q_l parameters and random effects $\mathbf{b} = [\mathbf{b}_1, \dots, \mathbf{b}_L] \sim N_{\sum q_l}(\mathbf{0}, \sigma_\varepsilon^2 \boldsymbol{\Omega}_b^{-1})$ and S smooth terms, we can write

$$\mathbf{y} = \tilde{\mathbf{X}}\tilde{\boldsymbol{\beta}} + \tilde{\mathbf{Z}}\tilde{\mathbf{b}} + \boldsymbol{\varepsilon}$$

with concatenated design matrices

$$\tilde{\mathbf{X}} = [\mathbf{X} \mathbf{X}_{u,1} \dots \mathbf{X}_{u,S}]; \quad \tilde{\mathbf{Z}} = [\mathbf{Z}_1 \dots \mathbf{Z}_L \mathbf{Z}_{p,1} \dots \mathbf{Z}_{p,S}]$$

and $\text{Cov}(\tilde{\mathbf{b}}) = \text{Cov} \left(\begin{bmatrix} \mathbf{b} \\ \mathbf{b}_1 \\ \vdots \\ \mathbf{b}_S \end{bmatrix} \right) = \sigma_\varepsilon^2 \begin{bmatrix} \boldsymbol{\Omega}_b^{-1} & 0 & \dots & 0 \\ 0 & \lambda_1 \mathbf{I}_{d_1 - p_1} & & \vdots \\ \vdots & & \ddots & 0 \\ 0 & \dots & 0 & \lambda_s \mathbf{I}_{d_S - p_S} \end{bmatrix}$

The parameter vector for the fixed effects is stacked in the same fashion as the one for the random effects. A minor additional complication arises from the fact that usually every matrix $\mathbf{X}_{u,s}$, $s = 1, \dots, S$ for the unpenalized parts of the smooth terms will contain an intercept column. They are removed in order to avoid a rank deficient design matrix $\tilde{\mathbf{X}}$. This also solves the identifiability issues common to additive models (Wood, 2006a, ch. 3.3).

2.1 Variability Estimation

The convenience functions `getF` and `plotF` to extract or plot estimated function values $\hat{f}(x)$ offer both MCMC-based or approximate frequentist variability bands. MCMC-based intervals (option `interval="MCMC"` in `getF`, `plotF`)

are pointwise HPD-intervals. They are based on samples from `lme4`'s `mcmc-samp` and may not be very reliable yet.¹

The frequentist variability estimates (option `interval="RW"` in `getF`, `plotF`) condition on the value of the estimated variance / smoothing parameters and use the bias-adjusted covariance of $\hat{f}(x)$ derived in Ruppert et al. (2003, ch. 6.4, eq. (6.13)). See section 3.5 for an example. I plan to include bootstrap-based variability estimates in future versions.

3 Examples

In this section, I fit some exemplary datasets to illustrate the capabilities of `amer`. I demonstrate how to fit simple semiparametric or additive models, how to use the `by`-option of the basis-generating function to fit group-specific smooths, how to use the `allPen`-option of the basis-generating function to fit subject- or cluster-specific smooth terms where all subject-level coefficients are penalized (i.e. the coefficients associated with \mathbf{X}_u are treated as random effects as well), and how to use the `varying`-option to fit varying-coefficient models. Most of the examples are adapted from Crainiceanu et al. (2005).

3.1 Generalized Additive Model

Let's first have a look at data on wages and union membership for 534 workers described in Berndt (1991). The model assumes that the probability of union membership of worker i ($y_i = 1$ if member) depends on his or her hourly wages x_i :

$$\begin{aligned} P(y_i = 1) &= \text{logit}^{-1}(f(x_i)) \\ f(x_i) &= \beta_0 + \beta_1 x_i + \sum_{k=1}^{K-1} b_k(x_i - \kappa_k)_+ \\ b_k &\sim N(0, \sigma_f^2) \end{aligned}$$

¹ Check the traceplots by calling `xyplot(attr(getF(<MyModel>, interval="MCMC"), "mcmc"))`, see section 3.5 for an example.

We use the default linear TP-Basis with the default number of basis function ($K = 15$) by calling the `tp`-function:

```
> u1 <- amer(UNION ~ tp(WAGE), family = binomial, data = union)
```

By default, `tp` uses equally spaced knots to generate the basis functions. If the covariate distribution is as non-uniform as here (see Figure 1), quantile-based knots are often a better choice:

```
> K <- 15
> degree <- 1
> knots <- quantile(union$WAGE, probs = (2:(K - degree +
+ 1))/(K - degree + 2))
> u2 <- amer(UNION ~ tp(WAGE, knots = knots), family = binomial,
+ data = union)

> print(u2, corr = F)
```

Generalized additive mixed model fit by the Laplace approximation

Formula: UNION ~ tp(WAGE, knots = knots)

Data: union

AIC	BIC	logLik	deviance
483	496	-238	477

Random effects:

Groups	Name	Variance	Std.Dev.
--------	------	----------	----------

f.WAGE	tp	0.202	0.449
--------	----	-------	-------

Number of obs: 534, groups: f.WAGE, 14

Fixed effects:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-0.961	0.727	-1.32	0.186
WAGE.fx1	1.837	0.890	2.06	0.039 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Note `amer`'s naming convention for the smooth function: The group name of the variance component associated with a smooth function of a covariate x is `f.x`, instead of the covariate name, `amer` gives the name of the basis generating function. The names of the columns in the $n \times p$ design matrix \mathbf{X}_u for the unpenalized part of the smooth are given by `x.fx1`, `x.fx2` to `x.fxp`. Figure 1 shows the plots produced by calls to `plotF`.

The following code generates figure 1:

```
> par(mfrow = c(1, 2))
> plotF(u1, trans = plogis, rug = F, ylim = c(0, 0.4),
+       auto.layout = F)
> with(union, points(WAGE, jitter(0.4 * UNION, factor = 0.15),
+                   cex = 0.5))
> plotF(u2, trans = plogis, rug = F, ylim = c(0, 0.4),
+       auto.layout = F)
> with(union, points(WAGE, jitter(0.4 * UNION, factor = 0.15),
+                   cex = 0.5))
```




Figure 1: Fitted probability of union membership versus hourly wages, with conditional 90% CI and jittered observations. Left panel: Equidistant-knots, right panel: quantile-based knots

3.2 Separate smooths for levels of a factor: Using the by-option

We use data on coronary sinus potassium concentration measurements for 36 dogs. The dogs were divided into 4 treatment groups, and the measurements for each dog were taken every two minutes from 1 to 13 minutes after occlusion (i.e. an artificially induced heart attack). The data was first published in Grizzle and Allen (1969) and previously analysed in Crainiceanu et al. (2005). Figure 2 shows the observed concentrations for all 36 dogs split up into the treatment groups. The group-averages seem to have quite different time trends, with different degrees of nonlinearity, so we fit an additive mixed model with group-specific smooth functions $f_g(t)$ ($g = 1, \dots, 4$) of time and random intercepts b_0 for the different dogs:

$$\begin{aligned}
 y_{ij} &= \beta_{0g(i)} + f_{g(i)}(t_{ij}) + b_{0i} + \varepsilon_{ij} \\
 f_{g(i)}(t_{ij}) &= \beta_{1g(i)} t_{ij} + \sum_{k=1}^{K-1} b_{f_{g(i)}k} (t_{ij} - \kappa_k)_+ \\
 b_{f_{g(i)}k} &\sim N(0, \sigma_{g(i)}^2) \\
 b_{0i} &\sim N(0, \sigma_{b_0}^2) \\
 \varepsilon_{ij} &\sim N(0, \sigma_\varepsilon^2)
 \end{aligned}$$

Note that we estimate separate spline coefficient variances σ_g^2 , $g = 1, \dots, 4$ for the 4 treatment groups.

The model is specified in `amer` using the `by`-option:

```

> data(dog)
> dl <- amer(y ~ -1 + group + tp(time, by = group) + (1 |
+      dog), data = dog)

> print(dl, corr = F)

Additive mixed model fit by REML
Formula: y ~ -1 + group + tp(time, by = group) + (1 | dog)
Data: dog
AIC BIC logLik deviance REMLdev
383 432 -177      342      355
Random effects:

```

Groups	Name	Variance	Std.Dev.
dog	(Intercept)	0.24870	0.499
f.time.group4	tp	0.00547	0.074
f.time.group3	tp	0.05706	0.239
f.time.group2	tp	0.00000	0.000
f.time.group1	tp	0.17814	0.422
Residual		0.15063	0.388

Number of obs: 252, groups: dog, 36; f.time.group4, 14; f.time.group3, 14; f.time.gr

Fixed effects:

	Estimate	Std. Error	t value
group1	4.3528	0.6008	7.24
group2	3.5529	0.1644	21.61
group3	4.3806	0.4380	10.00
group4	4.0446	0.2365	17.10
time.group1.fx1	0.1737	0.4230	0.41
time.group2.fx1	-0.0329	0.0465	-0.71
time.group3.fx1	0.5452	0.3080	1.77
time.group4.fx1	0.2285	0.1433	1.60

Note `amer`'s naming convention for smooth functions with a `by`-argument: The group name of the variance component associated with a smooth function of a covariate `x` at level `L` of the grouping factor `by` is `f.x.byL`. The names of the columns in the $n \times p$ design matrix $\mathbf{X}_{u,L}$ for the unpenalized part of the smooth for level `L` are given by `x.byL.fx1`, `x.byL.fx2` to `x.byL.fxp`.




Figure 2: `dog` data: coronary sinus potassium concentrations for 36 dogs in 4 treatment groups

The following code generates figure 3:

```
> layout(cbind(matrix(1, ncol = 2, nrow = 2), matrix(2:5,
+   ncol = 2, nrow = 2)))
> par(mar = c(3, 2.8, 2.8, 0.8), mgp = c(2, 1, 0))
> plotF(d1, ylim = range(dog$y), interval = "none", legend = "topleft",
+   level = 0.95, auto.layout = F, lwd = 3)
> dl.RW <- getF(d1, interval = "RW")
> for (i in 1:4) {
+   plot(0, 0, ylim = range(dog$y), xlim = c(0, 1), ylab = "y",
+     xlab = "time")
+   sub <- subset(dog, group == i)
+   lapply(split(sub, sub$dog, drop = T), function(x) lines(x$time,
+     x$y, col = "lightgrey", lty = 2, lwd = 0.5))
+   matlines(dl.RW[[1]][[i]][, 1], dl.RW[[1]][[i]][[,
+     -1], type = "l", lty = c(1, 3, 3), col = i, lwd = 2.5)
+ }
```




Figure 3: Left panel: Estimated groupwise smooths for the coronary sinus potassium data; Right panels: Estimated groupwise smooths with pointwise 90% CIs and observed data (grey dashes)

3.3 Subject- or cluster-specific smooths: Using the allPen-option

It is also possible to allow smooth subject-specific deviations from the group-specific curves. We fit this model with random intercepts and slopes for the dogs. The model is now

$$\begin{aligned}
y_{ij} &= \beta_{g(i)} + f_{g(i)}(t_{ij}) + f_i(t_{ij}) + \varepsilon_{ij} \\
f_{g(i)}(t_{ij}) &= \beta_{g(i)1}t_{ij} + \sum_{k=1}^K b_{f_{g(i)}k}(t_{ij} - \kappa_k)_+ \\
f_i(t_{ij}) &= b_{0i} + b_{1i}t_{ij} + \sum_{k=1}^K b_{f_ik}(t_{ij} - \kappa_k)_+ \\
b_{f_{g(i)}k} &\sim N(0, \sigma_{g(i)}^2) \\
b_{f_ik} &\sim N(0, \sigma_f^2) \\
(b_{0i}, b_{1i})' &\sim N_2(\mathbf{0}, \mathbf{D}) \\
\varepsilon_{ij} &\sim N(0, \sigma_\varepsilon^2)
\end{aligned}$$

We still estimate separate spline coefficient variances $\sigma_g^2, g = 1, \dots, 4$ for the 4 treatment groups, but only one common spline coefficient variance σ_f^2 for all the subject-specific smooth functions. We assume an unstructured covariance matrix \mathbf{D} for the subject-specific random intercepts and slopes (b_{0i}, b_{1i}) .

The model is specified in **amer** by using the **by**-option in combination with **allPen = TRUE**. To keep the dimension of the model reasonable, we use only **k= 5** basis functions per dog:

```

> d2 <- amer(y ~ -1 + group + tp(time, k = 5, by = dog,
+   allPen = T) + tp(time, by = group), data = dog)

> print(d2, corr = F)

```

Additive mixed model fit by REML

```

Formula: y ~ -1 + group + tp(time, k = 5, by = dog, allPen = T) + tp(time,
Data: dog
AIC BIC logLik deviance REMLdev
348 408 -157      301      314

```

by =

Random effects:

Groups	Name	Variance	Std.Dev.	Corr
f.time.dog	tp	0.04051	0.2013	
u.time.dog	(Intercept)	0.25259	0.5026	
	time.dog.fx1	0.00468	0.0684	1.000
f.time.group4	tp	0.00587	0.0766	
f.time.group3	tp	0.05887	0.2426	
f.time.group2	tp	0.00000	0.0000	
f.time.group1	tp	0.20025	0.4475	
Residual		0.09588	0.3096	

Number of obs: 252, groups: f.time.dog, 144; u.time.dog, 36; f.time.group4, 14; f.time.group3, 14; f.time.group2, 14; f.time.group1, 14

Fixed effects:

	Estimate	Std. Error	t value
group1	4.3168	0.5797	7.45
group2	3.5742	0.1774	20.15
group3	4.3174	0.4179	10.33
group4	4.0878	0.2389	17.11
time.group1.fx1	0.1453	0.4000	0.36
time.group2.fx1	-0.0115	0.0773	-0.15
time.group3.fx1	0.5039	0.2861	1.76
time.group4.fx1	0.2715	0.1436	1.89

By specifying `allPen = TRUE`, a random intercept for the `by`-variable is automatically included in the model. Also note `amer`'s naming convention for smooth functions of a covariate `x` with a `by`-argument and `allPen = TRUE`: For the random effects associated with \mathbf{X}_u , the group name of the variance component is `u.x.by`. The factor `u.x.by` is of course the same as `by`, the renaming is done for technical reasons.

Especially for spline bases with a higher dimensional nullspace of the penalty it may not be feasible or desirable to estimate an unstructured covariance matrix \mathbf{D} . By setting the `diag`-option to `TRUE` in the specification of a smooth term with `allPen = TRUE`, we can enforce uncorrelated random effects for the coefficients associated with \mathbf{X}_u :

```
> d3 <- amer(y ~ -1 + group + tp(time, k = 5, by = dog,
+   allPen = T, diag = T) + tp(time, k = 5, by = group),
+   data = dog)

> print(d3, corr = F)
```



```

Additive mixed model fit by REML
Formula: y ~ -1 + group + tp(time, k = 5, by = dog, allPen = T, diag = T) +      tp(
  Data: dog
      AIC BIC logLik deviance REMLdev
    348 405   -158     301     316
Random effects:
      Groups      Name      Variance Std.Dev.
f.time.dog      tp      0.0409    0.202
u.time.dog      time.dog.fx1 0.0000    0.000
u.time.dog      (Intercept) 0.2000    0.447
f.time.group4    tp      0.0176    0.133
f.time.group3    tp      0.2017    0.449
f.time.group2    tp      0.0000    0.000
f.time.group1    tp      0.6588    0.812
Residual                        0.0970    0.311
Number of obs: 252, groups: f.time.dog, 144; u.time.dog, 36; f.time.group4, 4; f.time.group3, 4; f.time.group2, 4; f.time.group1, 4

Fixed effects:
      Estimate Std. Error t value
group1      4.3666    0.3375   12.94
group2      3.5742    0.1621   22.05
group3      4.3521    0.3029   14.37
group4      4.0829    0.2039   20.03
time.group1.fx1 0.1769    0.2448    0.72
time.group2.fx1 -0.0116    0.0746   -0.16
time.group3.fx1 0.5256    0.2148    2.45
time.group4.fx1 0.2676    0.1247    2.15

```

3.4 Varying coefficient models: Using the varying-option

Another class of models that can be fitted with `amer` are varying coefficient models. They are used to model smoothly varying regression coefficients, i.e. models in which the effect of a covariate z varies smoothly over the range of another covariate x (\cdot denotes elementwise multiplication.):

$$\begin{aligned} \mathbf{y} &= \beta(\mathbf{x}) \cdot \mathbf{z} + \varepsilon \\ \beta(\mathbf{x}) &= f(\mathbf{x}) \approx \mathbf{X}_u \boldsymbol{\beta} + \mathbf{Z}_p \mathbf{b} \\ \Rightarrow \beta(\mathbf{x}) \cdot \mathbf{z} &\approx (\mathbf{X}_u \cdot \mathbf{z}) \boldsymbol{\beta} + (\mathbf{Z}_p \cdot \mathbf{z}) \mathbf{b} \end{aligned}$$

This class of models can be fitted by simply scaling the design matrices for the spline of the effect-modifying covariate x (i.e. the varying coefficient) with the values of the covariate z . A slight complication arises: for all other classes of models, we drop the intercept column in \mathbf{X}_u so that the model is identifiable. That is unnecessary in this case, so the design matrix $(\mathbf{X}_u \cdot \mathbf{z})$ has $\mathbf{1} \cdot \mathbf{z} = \mathbf{z}$ as its first column.

Let's look at `lattice`'s `ethanol` data set as an example: Ethanol fuel was burned in a single-cylinder engine. For various settings of the engine compression (`C`) and the equivalence ratio (`E`, a measure of the richness of the air and ethanol fuel mixture), the emissions of nitrogen oxides (`NOx`) were recorded. We assume that, for a given equivalence ratio, the relationship between compression and emissions is linear with different intercepts (see figure 4).

The model we want to fit is

$$\begin{aligned} \text{NOx}_i &= f_1(\text{E}_i) + f_2(\text{E}_i)\text{C}_i + \varepsilon_i \\ f_1(\text{E}_i) &= \beta_0 + \mathbf{X}_u^{(E)} \boldsymbol{\beta}^{(E)} + \mathbf{Z}_p^{(E)} \mathbf{b}^{(E)} \\ f_2(\text{E}_i)\text{C}_i &= \mathbf{X}_u^{(EC)} \boldsymbol{\beta}^{(EC)} + \mathbf{Z}_p^{(EC)} \mathbf{b}^{(EC)}, \end{aligned}$$

with the usual distributional assumptions about $\mathbf{b}^{(E)}, \mathbf{b}^{(EC)}$ and ε . The command to fit this model in `amer` is simply

```
> e1 <- amer(NOx ~ tp(E, k = 20) + tp(E, k = 20, varying = C),
+           data = ethanol)

> print(e1, corr = F)
```




Figure 4: Emissions of nitrogen oxides NO_x for various engine compression values C , split up according to equivalence ratio E . Lines are linear regression estimates for the subgroups.

```

Additive mixed model fit by REML
Formula: NOx ~ tp(E, k = 20) + tp(E, k = 20, varying = C)
Data: ethanol
AIC BIC logLik deviance REMLdev
15.9 33.2 -0.936 -10.2 1.87
Random effects:
Groups Name Variance Std.Dev.
f.E tp 1.286220 1.1341
f.EXC tp 0.000988 0.0314
Residual 0.030132 0.1736
Number of obs: 88, groups: f.E, 19; f.EXC, 19

Fixed effects:
Estimate Std. Error t value
(Intercept) 2.1807 2.0905 1.043
E.fx1 1.6448 1.3032 1.262
EXC.fx1 0.1324 0.0844 1.568
EXC.fx2 0.0273 0.0550 0.497

```

The fit is plotted in figure 5. Note `amer`'s naming convention for varying coefficient models: For an effect-modifying covariate x and an effect-causing covariate z , the function name is given as `f.xXz`, the unpenalized effects are named `xXz.fx1` to `xXz.fxp`. The first unpenalized effect corresponds to the conventional regression coefficient for z , since \mathbf{X}_u has z as its first column.

3.5 Variability Bands: Using the RW- and MCMC-options

By default, `amer`'s `plotF` requests and plots approximate pointwise intervals based on a bias-adjusted approximation of $\text{Cov}((\hat{\beta}, \hat{\mathbf{b}} - \mathbf{b}) | \hat{\sigma}_b^2, \hat{\sigma}_\varepsilon^2)$ (see 2.1). These may underestimate the true variability since they ignore the uncertainty in the estimated variance parameters. We can also use the results from `lme4`'s `mcmc` to construct MCMC-based variability bands. Figure 5 compares the frequentist, bias-adjusted variability bands (see 2.1) for the estimated function values with pointwise HPD-Intervals based on 1000 draws from `mcmc` for the `ethanol` data. The latter are much wider since they take into account the variability of the estimated variances, while the former condition on the estimated variances.

Since `mcmc` may not always work as expected, it is strongly recommended to examine the returned MCMC-samples. They are available as the `mcmc`-attribute of the value returned by `getF` or `plotF`. A quick visual inspection of the sampling paths can be done via `xyplot`, see figure 6. The MCMC iterations show strange spikes after about 700 iterations with humongous values drawn for the relative standard deviations `e1@ST` of the spline coefficients. The marginal posterior densities for the variance components are concentrated on values magnitudes larger than the REML estimates found by the optimizer, with ridiculously long upper tails (which lead to erratic sampling behaviour of the spline coefficients \mathbf{b} and consequently enormous HPD-Intervals for $\hat{f}(\mathbf{x})$):

```
> e1.MCMCData <- as.data.frame(attr(e1.MCMC, "mcmc"))
> data.frame(c(fixef(e1), e1@ST, lme4:::sigma(e1)))
```

	X.Intercept.	E.fx1	EXC.fx1	EXC.fx2	tp	tp.1	sigmaREML
tp	2.18	1.64	0.132	0.0273	6.53	0.181	0.174

```
> apply(e1.MCMCData, 2, quantile, probs = c(0.1, 0.25,
+      0.5, 0.75, 0.9), na.rm = T)
```

	(Intercept)	E.fx1	EXC.fx1	EXC.fx2	ST1	ST2	sigma
10%	-48.15	-28.17	-4.931	-2.778	0.00e+00	3.05e+01	0.123
25%	-9.27	-6.17	-2.985	-1.640	6.28e+03	2.12e+08	0.130
50%	7.96	4.20	-0.629	-0.317	4.20e+10	6.36e+24	0.139
75%	49.78	27.92	0.354	0.231	4.46e+21	8.89e+52	0.154
90%	85.92	47.91	2.399	1.390	3.12e+35	1.22e+83	0.243

The following code generates figure 5:

```
> par(mfrow = c(2, 2), mar = c(3, 2.8, 2.8, 0.8), mgp = c(2,
+   1, 0))
> el.RW <- plotF(el, addConst = c(T, F), level = 0.95,
+   auto.layout = F)
> set.seed(12345)
> el.MCMC <- plotF(el, addConst = c(T, F), int = "MCMC",
+   sims = 1000, level = 0.95, auto.layout = F)
```

starting 1000 MCMC iterations for posterior intervals:
... done.




Figure 5: Fits and 95% CIs/HPD-Intervals for the `ethanol` data. Upper row: pointwise frequentist variability bands conditional on the estimated variances. Lower row: pointwise HPD-Intervals based on 1000 draws (see text) from `mcmcscamp`. Left column: Effect of equivalence ratio `E`. Right column: regression coefficient for compression `C` varying over `E`. By default, the value of the varying coefficient function²² (right column) is evaluated for a co-variate value of $z = 1$, so the plot for `f.EXC` can be interpreted directly as $\beta(E)$.

⇒ we cannot blindly trust MCMC-intervals until `mcmc` becomes more stable.

```
> print(xyplot(attr(e1.MCMC, "mcmc")))
```




Figure 6: Traceplot of the MCMC samples for model `e1`. Note the huge values for `ST`.

4 Implementation

4.1 What's the point?

There is already a well-tested, well documented and versatile package `mgcv` (Wood, 2006b) that fits generalized additive mixed models in R, so why bother with yet another one?

- `mgcv`'s `gamm` uses the less stable and slower `nlme`-implementation of linear mixed models, while `amer` relies on the more stable algorithm used in `lme4` (Bates and Maechler, 2009b) with its very fast sparse `Matrix` (Bates and Maechler, 2009a) magic.
- `mgcv`'s `gamm` fits non-gaussian responses by calling MASS's `glmmPQL`. The PQL-approach for fitting generalized linear mixed models (GLMM) is severely biased and often unstable. `amer` relies on the more precise Laplace approximation implemented in `lme4` for fitting GLMMs.
- `asreml` also offers additive models, but is limited to gaussian responses (and isn't free or open-source)

The drawbacks of using `amer` instead of `mgcv`'s `gamm` are that, as yet, it's not possible to include serial and/or spatial correlation structures or variance functions for the residuals or specify covariance structures of the random effects that aren't either diagonal or unstructured (this will remain an issue as long as `lme4` doesn't have that capability). Also, multidimensional smooths are not yet implemented, but will hopefully be included in a future version.

4.2 Making `lmer` fit GAMMs

In its most current version (0.999375-31, at the time of writing), `lme4` fits mixed models

- for hierachical data structures (i.e. grouped data) and
- only admits either diagonal or unstructured covariances of the non-scalar random effects for every level of grouping.

In an additive mixed model, data are not grouped (the smoothing introduces dependence between all the observation in the data) and, if the reparametrization from $\mathbf{B}\boldsymbol{\delta}$ to $\mathbf{X}_u\boldsymbol{\beta} + \mathbf{Z}_p\mathbf{b}$ is not done, the precision matrix \mathbf{K}/λ for the penalized coefficients is in general not diagonal (and certainly not unstructured) and not of full rank, so that an implementation of GAMMs based on the unreparameterized representation is not possible without changing the underlying C-code of `lme4`.

Instead of making these changes to the underlying C, `amer` tricks `lmer` into fitting additive models by setting up an unfit model object with the structure of random and fixed effect design matrices necessary for the mixed model representation (2) of the additive model, and then overwriting the (precursor of) the `Zt`-slots with the penalized parts \mathbf{Z}_p of the reparameterized spline bases. More precisely, the model object is set up by going through the following steps for each smooth function:

1. Generate \mathbf{X}_u and \mathbf{Z}_p according to the basis generating function (see section 4.4) given for the smooth term,
2. replace smooth term in the original model formula with fixed effect terms for the columns in \mathbf{X}_u and a random intercept term for an artificial grouping factor that has as many levels as \mathbf{Z}_p has columns,
3. add the fixed effects in \mathbf{X}_u and the artificial grouping factor to the model frame,
4. set up, but do not fit this model with a call to `lmer` with option `doFit=FALSE`, and finally
5. overwrite the design matrices for the random intercept of the artificial grouping factor with \mathbf{Z}_p .

Some complications arise if the `by-` or `allPen`-option are used, but these steps remain basically the same. The routine then hands back the modified unfitted model to `lmer_finalize` or `glmer_finalize` for calling the optimization C-code etc.

4.3 Justification for using the TP-Basis

The TP-basis implemented in `amer` suffers from some serious flaws:

- It has an undesirable one-to-one mapping between the smoothness / differentiability of the fit (TP of degree $p \Rightarrow$ fit is $p - 1$ -time continuous differentiable), and the nullspace of the penalty (TP of degree $p \Rightarrow$ nullspace is a p -degree polynomial). This is different from, e.g., penalized B-Spline fits, where the order of the difference penalty that determines the nullspace of the penalty can be specified independently from the order of the spline bases which determine the smoothness of the fit.
- the unbounded support (to the right of the knot) of the truncated polynomials means that the values of the basis function can potentially become huge.
- the columns in \mathbf{Z}_p containing the truncated polynomials are severely collinear, especially for closely spaced knots

However, similar collinearity in \mathbf{Z}_p is present for all other spline bases after the mixed model reparametrization (2) described in section 2. This not well-known fact and the fact that for all other spline bases I am aware of, the reparameterized basis in \mathbf{Z}_p is no longer sparse, even if the original \mathbf{B} was are the reasons why I chose the TP-Basis as the default for `amer`: Since \mathbf{Z}_p is about 50% zeroes, `amer` can take full advantage of the sparse matrix operations used in `lme4`.

4.4 Writing your own basis-generating function

It is fairly easy to implement your own basis generating function for use in `amer`. Such a function only has to fulfill the following criteria:

- It has to have arguments
 - `x`, a `numeric` variable used for the smooth function,
 - `by`, a `factor` variable (default: `NULL`),
 - `allPen`, (a `logical`),
 - `diag`, (a `logical`),
 - `varying`, a `numeric` variable (default: `NULL`).

- It has to return a list with
 - an entry named **X**, which contains the matrix X_u without the intercept column (this can be a matrix with zero columns)
 - an entry named **Z**, which contains the matrix Z_p
 - an attribute call, which contains the result of `expand.call`

The technical details of splitting up X_u and Z_p for a possible `by` variable, naming the columns in X_u etc. are performed by the utility function `expandBasis`.

As an example, let's add a variant of the TP-Basis to `amer`'s repertoire – let's say we want to get rid of the undesirable one-to-one mapping between the smoothness/differentiability of the fit and the null-space of the penalty of the TP-Basis. We implement a simple basis-generating function `tp2` that lets us specify the dimensionality of the nullspace so that, for a TP-spline basis of degree p without intercept (see above), we can specify the degree of the global polynomial that is unpenalized. Let's call this option `dimU`. If we let `dimU = p`, this corresponds to the conventional TP-Penalty. If `dimU < p`, columns containing global polynomials that would be in X_u for the conventional TP-Penalty are put in Z_p instead.

The following code implements a rough draft of the idea, with the default for using a quadratic TP-Basis (`p=2`) (s.t. the fitted function is continuously differentiable, i.e. has no kinks) while penalizing deviations from linearity (`dimU=1`):

```
> tp2 <- function(x, p = 2, k = 15, dimU = 1, by = NULL,
+               allPen = FALSE, diag = FALSE, varying = NULL,
+               knots = quantile(x,
+                               probs = (2:(k - p + 1))/(k - p + 2)))
+ {
+   #dim. of nullspace can't be larger than p of TP-basis:
+   stopifnot(dimU <= p)
+
+   #always need this for the call attribute of the returned value:
+   call <- as.list(expand.call())
+   call$knots <- knots
+
+   #global polynomial trends (no intercept!):
+   X <- if (p > 0) {
+     outer(x, 1:p, "^")
+   } else {
+     matrix(nrow = length(x), ncol = 0)
```

```

+           }
+
+       #TP-design for penalised part:
+       Z <- outer(x, knots, "-")^p * outer(x, knots, ">")
+
+       # adapt design for dimU option:
+       if (dimU != p) {
+           Xp <- X[, (1:p) > dimU, drop = F]
+           X <- X[, (1:p) <= dimU, drop = F]
+           Z <- cbind(Xp, Z)
+       }
+
+       res <- list(X = X, Z = Z)
+       attr(res, "call") <- as.call(call)
+       return(res)
+ }

```

We can now use this function to fit a continuously differentiable function with penalized deviations from linearity to the dog data, but we have to tell `amer` to look for smooth terms called `tp2` in the `basisGenerators`-option:

```

> d4 <- amer(y ~ -1 + group + tp2(time, k = 5, p = 2, dimU = 1,
+   by = group) + (1 | dog), data = dog, basisGenerators = c("tp2"))

> print(d4, corr = F)

```

Additive mixed model fit by REML

Formula: `y ~ -1 + group + tp2(time, k = 5, p = 2, dimU = 1, by = group) + (1 | dog)`
 Data: dog

AIC	BIC	logLik	deviance	REMLdev
374	423	-173	340	346

Random effects:

Groups	Name	Variance	Std.Dev.
dog	(Intercept)	0.249	0.499
f.time.group4	tp2	0.173	0.416
f.time.group3	tp2	2.096	1.448
f.time.group2	tp2	0.000	0.000
f.time.group1	tp2	34.867	5.905
Residual		0.150	0.387

Number of obs: 252, groups: dog, 36; f.time.group4, 4; f.time.group3, 4; f.time.group2, 4; f.time.group1, 4

Fixed effects:

	Estimate	Std. Error	t value
group1	4.0891	0.2060	19.85

group2	3.6021	0.1785	20.18
group3	3.5420	0.2117	16.73
group4	3.7040	0.1941	19.08
time.group1.fx1	0.1636	1.1119	0.15
time.group2.fx1	-0.0986	0.1389	-0.71
time.group3.fx1	2.1733	0.6108	3.56
time.group4.fx1	0.6959	0.3559	1.96

Figure 7 shows a comparison of the fit with 5 basis functions of the `tp2`-function to the fit of a conventional linear TP-basis.

The following code generates figure 7:

```
> d1.k5 <- amer(y ~ -1 + group + tp(time, k = 5, by = group) +
+ (1 | dog), data = dog)
> par(mfrow = c(1, 2))
> plotF(d1.k5, legend = "topleft", auto.layout = F)
> plotF(d4, legend = "none", auto.layout = F)
```




Figure 7: Comparison of the results for `tp(degree=1)` (left panel) and `tp2(degree=2, dimU=1)` (right panel) for the dog data.

5 Open Issues

A to-do list for developping `amer` further:

- approximate frequentist CI's for smooths with `allPen=TRUE` (should be easy, only modify `fctV`)
- 2D-smooths (will mean major reworking of most utility functions called by `amerSetup` as well as `getF/PlotF`)
- implementing (parametric/wild/...) bootstrap-CIs (could use `lme4::refit` and Ben Bolker's `mer.sim`, maybe implement Kauermann/Claeskens/Opsomer(2008))
- implementing other spline bases, e.g. for cyclic functions

References

- Douglas Bates and Martin Maechler. *Matrix: Sparse and Dense Matrix Classes and Methods*, 2009a. URL <http://CRAN.R-project.org/package=Matrix>. R package version 0.999375-30.
- Douglas Bates and Martin Maechler. *lme4: Linear mixed-effects models using Eigen and Eigenpack*, 2009b. URL <http://CRAN.R-project.org/package=lme4>. R package version 0.999375-31.
- E. Berndt. *The Practice of Econometrics: Classical and Contemporary*. Addison - Wesley, 1991.
- C. Crainiceanu, D. Ruppert, and M.P. Wand. Bayesian analysis for penalized spline regression using WinBUGS. *Journal of Statistical Software*, 14(14):1–24, 2005.
- P.H.C. Eilers and B.D. Marx. Flexible smoothing with B-splines and penalties. *Statistical Science*, 11(2):89–121, 1996.
- J.E. Grizzle and D.M. Allen. Analysis of growth and dose response curves. *Biometrics*, 25:357–381, 1969.
- T. Kneib. *Mixed model based inference in structured additive regression*. PhD thesis, LMU München, 2006.

- D. Ruppert, M.P. Wand, and R.J. Carroll. *Semiparametric Regression*. Cambridge University Press, 2003.
- S. Wood. *Generalized Additive Models*. Chapman and Hall, 2006a.
- S. Wood. *mgcv: Multiple Smoothing Parameter Estimation by GCV or UBRE*, 2006b. URL <http://www.maths.bath.ac.uk/~sw283/>.