

flexsurv: flexible parametric survival modelling in R. Supplementary examples

Christopher H. Jackson
MRC Biostatistics Unit, Cambridge, UK
chris.jackson@mrc-bsu.cam.ac.uk

Abstract

This vignette of examples supplements the main **flexsurv** user guide.

Keywords: ~survival.

1. Examples of custom distributions

1.1. Proportional hazards generalized gamma model

Crowther and Lambert (2013) discuss using the **stgenreg** Stata package to construct a proportional hazards parameterisation of the three-parameter generalised gamma distribution. A similar trick can be used in **flexsurv**. A four-parameter custom distribution is created by defining its hazard (and cumulative hazard) functions. These are obtained by multiplying the built-in functions **hgengamma** and **Hgengamma** by an extra dummy parameter, which is used as the location parameter of the new distribution. The intercept of this parameter is fixed at 1 when calling **flexsurvreg**, so that the new model is no more complex than the generalized gamma AFT model **fs3**, but covariate effects on the dummy parameter are now interpreted as hazard ratios.

```
R> library(flexsurv)
R> hgengammaPH <- function(x, dummy, mu=0, sigma=1, Q){
+   dummy * hgengamma(x=x, mu=mu, sigma=sigma, Q=Q)
+ }
R> HgengammaPH <- function(x, dummy, mu=0, sigma=1, Q){
+   dummy * Hgengamma(x=x, mu=mu, sigma=sigma, Q=Q)
+ }
R> custom.gengammaPH <- list(name="gengammaPH",
+   pars=c("dummy","mu","sigma","Q"), location="dummy",
+   transforms=c(log, identity, log, identity),
+   inv.transforms=c(exp, identity, exp, identity),
+   inits=function(t){
+     lt <- log(t[t>0])
+     c(1, mean(lt), sd(lt), 0)
```

```

+                               })
R> fs7 <- flexsurvreg(Surv(recyrs, censrec) ~ group, data=bc,
+                   dist=custom.gengammaPH, fixedpars=1)

```

2. Examples of custom model summaries

2.1. Plotting a hazard ratio against time

The following code plots the hazard ratio (Medium versus Good prognostic group) against time for both the proportional hazards model `fs7` and the better-fitting accelerated failure time model `fs2`. It illustrates the use of the following functions.

`summary.flexsurvreg` for generating the estimated hazard at a series of times, for particular covariate categories.

`normboot.flexsurvreg` for generating a bootstrap-style sample from the sampling distribution of the parameter estimates, for particular covariate categories.

`do.call` for constructing a function call by supplying a list containing the function's arguments. This is used throughout the source of **flexsurv**.

```


R> fs2 <- flexsurvreg(Surv(recyrs, censrec) ~ group + sigma(group),
+                   data=bc, dist="gengamma")
R> B <- 5000
R> t <- seq(0.1, 8, by=0.1)
R> hrAFT.est <-
+   summary.flexsurvreg(fs2, t=t, type="hazard",
+                       newdata=data.frame(group="Medium"), ci=FALSE)[[1]][, "est"] /
+   summary.flexsurvreg(fs2, t=t, type="hazard",
+                       newdata=data.frame(group="Good"), ci=FALSE)[[1]][, "est"]
R> pars <- normboot.flexsurvreg(fs2, B=B, newdata=data.frame(group=c("Good", "Medium")))
R> hrAFT <- matrix(nrow=B, ncol=length(t))
R> for (i in seq_along(t)){
+   haz.medium.rep <- do.call(hgengamma, c(list(t[i]), as.data.frame(pars[[2]])))
+   haz.good.rep <- do.call(hgengamma, c(list(t[i]), as.data.frame(pars[[1]])))
+   hrAFT[,i] <- haz.medium.rep / haz.good.rep
+ }
R> hrAFT <- apply(hrAFT, 2, quantile, c(0.025, 0.975))
R> hrPH.est <-
+   summary.flexsurvreg(fs7, t=t, type="hazard",
+                       newdata=data.frame(group="Medium"), ci=FALSE)[[1]][, "est"] /
+   summary.flexsurvreg(fs7, t=t, type="hazard",
+                       newdata=data.frame(group="Good"), ci=FALSE)[[1]][, "est"]
R> pars <- normboot.flexsurvreg(fs7, B=B, newdata=data.frame(group=c("Good", "Medium")))
R> hrPH <- matrix(nrow=B, ncol=length(t))

```

```

R> for (i in seq_along(t)){
+   haz.medium.rep <- do.call(hgengammaPH, c(list(t[i]), as.data.frame(pars[[2]])))
+   haz.good.rep <- do.call(hgengammaPH, c(list(t[i]), as.data.frame(pars[[1]])))
+   hrPH[,i] <- haz.medium.rep / haz.good.rep
+ }
R> hrPH <- apply(hrPH, 2, quantile, c(0.025, 0.975))
R> plot(t, hrAFT[1,], type="l", ylim=c(0, 10), col="red", xlab="Years",
+   ylab="Hazard ratio (Medium / Good)", lwd=1, lty=2)
R> lines(t, hrAFT[2,], col="red", lwd=1, lty=2)
R> lines(t, hrPH[1,], col="darkgray", lwd=1, lty=2)
R> lines(t, hrPH[2,], col="darkgray", lwd=1, lty=2)
R> lines(t, hrAFT.est, col="red", lwd=2)
R> lines(t, hrPH.est, col="darkgray", lwd=2)
R> legend("topright", lwd=c(2,2), col=c("red","darkgray"), bty="n",
+   c("Generalized gamma: standard AFT", "Generalized gamma: proportional hazards"))

```



2.2. Restricted mean survival

The expected survival up to time t , from a model with cumulative distribution $F(t|\alpha)$, is

$$E(T|T < t) = \int_0^t 1 - F(u|\alpha) du$$

An estimate and confidence interval for this, for a specified covariate value, can be computed using a custom summary function as follows. (Note that `summary.flexsurvreg` can be abbreviated to `summary`). As in the `median.weibull` example in the user guide vignette, the summary function is independent of time, so any value can be specified for `t` in the call to `summary`. The time horizon up to which to compute the mean is specified by the default value of the `horizon` argument to the custom function. The mean survival is computed here up to 100 years. Setting `horizon=Inf` is theoretically also possible for an unrestricted mean, but the integral does not converge in this example.

```
R> mean.gengamma <- function(mu, sigma, Q, horizon=100, ...){
+   surv <- function(t, ...) { 1 - pgengamma(q=t, mu=mu, sigma=sigma, Q=Q, ...) }
+   integrate(surv, 0, horizon, ...)$value
+ }
R> summary(fs2, newdata=list(group="Good"), t=1, fn=mean.gengamma)
```

```
group=Good
  time      est      lcl      ucl
1    1 21.50102 13.72859 32.58824
```

```
R> summary(fs2, newdata=list(group="Medium"), t=1, fn=mean.gengamma)
```

```
group=Medium
  time      est      lcl      ucl
1    1 12.00025  8.028092 17.96442
```

```
R> summary(fs2, newdata=list(group="Poor"), t=1, fn=mean.gengamma)
```

```
group=Poor
  time      est      lcl      ucl
1    1  5.539196  3.808509  8.97207
```

Note that the (unrestricted) median is more stable, and less than the restricted mean due to the skewness of this distribution.

```
R> median.gengamma <- function(mu, sigma, Q) {
+   qgengamma(0.5, mu=mu, sigma=sigma, Q=Q)
+ }
R> summary(fs2, newdata=list(group="Good"), t=1, fn=median.gengamma)
```

```
group=Good
  time      est      lcl      ucl
1    1  9.736555  7.171817 13.60485
```

3. Spline models

3.1. Prognostic model for the German breast cancer data

The regression model III in [Sauerbrei and Royston \(1999\)](#) used to create the prognostic group from the breast cancer data (supplied as `bc` in `flexsurv` and `GBSG2` in `TH.data`) can be reproduced as follows. Firstly, the required fractional polynomial transformations of the covariates are constructed. `progc` implements the Cox model used by [Sauerbrei and Royston \(1999\)](#), and `prog3` is a flexible fully-parametric alternative, implemented as a spline with three internal knots. The number of knots was chosen to minimise AIC. The covariate effects are very similar.

After fitting the model, the prognostic index can then be derived from categorising observations in three groups according to the tertiles of the linear predictor in each model. The indices produced by the Cox model (`progc`) and the spline-based model (`progf`) agree exactly.

```
R> if (require("TH.data")){
+
+   GBSG2 <- transform(GBSG2,
+                       X1a=(age/50)^-2,
+                       X1b=(age/50)^-0.5,
+                       X4=tgrade %in% c("II","III"),
+                       X5=exp(-0.12*pnodes),
+                       X6=(progrecc+1)^0.5
+                       )
+   (progc <- coxph(Surv(time, cens) ~ horTh + X1a + X1b + X4 +
+                   X5 + X6, data=GBSG2))
+   (prog3 <- flexsurvspline(Surv(time, cens) ~ horTh + X1a + X1b + X4 +
+                             X5 + X6, k=3, data=GBSG2))
+   predc <- predict(progc, type="lp")
+   progc <- cut(predc, quantile(predc, 0:3/3))
+   predf <- model.matrix(prog3) %*% prog3$res[-(1:5),"est"]
+   progf <- cut(predf, quantile(predf, 0:3/3))
+   table(progc, progf)
+ }
```

	progf		
progc	(-9.91,-7.76]	(-7.76,-7.11]	(-7.11,-3.36]
(-2.52,-0.359]	228	0	0
(-0.359,0.285]	0	228	0
(0.285,4]	0	0	229

References

- Crowther MJ, Lambert PC (2013). “**stgenreg**: A Stata Package for General Parametric Survival Analysis.” *Journal of Statistical Software*, **53**, 1–17.
- Sauerbrei W, Royston P (1999). “Building Multivariable Prognostic and Diagnostic Models: Transformation of the Predictors by Using Fractional Polynomials.” *Journal of the Royal Statistical Society A*, **162**(1), 71–94.